

Intestinal Parasitic Infections in Man: A Review

M Norhayati, PhD, M S Fatmah, Dip Med Tech, S Yusof, Dip Med Tech, A B Edariah, MPHTM

Department of Parasitology and Medical Entomology, Faculty of Medicine, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur

Summary

Intestinal parasites of importance to man are *Enterobius vermicularis*, the soil-transmitted helminthes (STH) - *Ascaris lumbricoides*, *Trichuris trichiura*, hookworms (*Necator americanus*/*Ancylostoma duodenale*) and *Strongyloides stercoralis* and the protozoa *Entamoeba histolytica* and *Giardia duodenalis*. Other protozoa such as *Cryptosporidium* sp. and *Isospora* sp. are becoming important in causing prolonged diarrhea in immunocompromised patients. It is estimated that almost 1 billion, 500 million and 900 million people worldwide are infected by the major nematode species - *A. lumbricoides*, *T. trichiura* and hookworms respectively. Most of the infections are endemic and widely distributed throughout poor and socio-economically deprived communities in the tropics and subtropics. Environmental, socio-economic, demographic and health-related behavior is known to influence the transmission and distribution of these infections. In giardiasis, one study indicates that age ≤ 12 years and the presence of family members infected with *Giardia* were risk factors for infection. Most of the infections occur in children and both genders are equally affected. Epidemiological studies of STH infections have shown that the prevalence and intensity of infection are highest among children 4-15 years of age. The frequency of distribution of STH infections is over-dispersed and highly aggregated. Predisposition to reinfection following treatment has been reported in these infections. In highly endemic areas reinfection can occur as early as 2 months post-treatment, and by 4 months, almost half of the population treated become reinfected. By 6 months the intensity of infection was similar to pretreatment level.

Key Words: Intestinal Parasitic Infections

Introduction

Intestinal parasitic infections are distributed throughout the world, with high prevalence in poor and socio-economically deprived communities in the tropics and subtropics. Amebiasis, ascariasis, hookworm infection and trichuriasis are among the top ten most common infections in the world¹. These infections continue

to be a global health problems, particularly among children in poor communities in developing countries². If AIDS is synonymous with promiscuity, intestinal parasitic infections are synonymous with poverty. Epidemiological assessments of these infections have traditionally focused on estimating the number of infections (prevalence), which occur worldwide. This is

This article was accepted: 22 September 2002

Corresponding Author: Norhayati Moktar, Department of Parasitology, Faculty of Medicine, University Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur.

because the clinical consequences of these infections are very mild.

The Global Burden of Intestinal Parasitic Infections

Intestinal Helminthic Infections

Intestinal helminths of importance to man are *E. vermicularis* (pinworm), soil-transmitted helminths (STH) - *A. lumbricoides* (round worm), *T. trichiura* (whip worm), *N. americanus* and *A. duodenale* (hookworm) and *S. stercoralis* (threadworm). The other intestinal nematodes (*Anisakis* sp., *Capillaria philippinensis*), trematodes and cestodes are less widespread in man. Their distribution is limited to certain areas in the world and the infections are usually confined to certain communities.

It is estimated that 25% of the world population are infected by *A. lumbricoides* and this causes up to a million cases of disease annually^{3,4}; 500-600 million people worldwide are infected by *T. trichiura*⁵ and about 500 million by hookworm⁶. The distribution of *S. stercoralis* infection usually follows that of hookworm. It is estimated that 50-100 million of the world's population are infected by *S. stercoralis*. The worldwide infection by *Enterobius vermicularis* is about 200 million and it is the commonest helminthic infection in the United States (40 million). In contrast to soil-transmitted helminthiasis, enterobiasis is prevalent in both developed and developing countries^{7,8}. Intestinal helminthic infections are endemic and widely distributed throughout poor and socio-economically deprived communities in the tropics and subtropics; where poverty, overcrowding, poor environmental sanitation and low level of education are more apparent problems than intestinal helminthic infections (Table I).

Intestinal Protozoal Infections

Intestinal protozoa of importance to man are *Entamoeba histolytica* and *Giardia duodenalis*.

Currently *Blastocystis hominis* and opportunistic protozoa such as *Cryptosporidium* sp. and *Isospora* sp. have been identified as the causes of diarrhea in children and immunocompromised patients. Other protozoal intestinal infections have restricted distribution (*Balantidium coli*) or are widely distributed but not pathogenic (*Entamoeba coli*, *Dientamoeba fragilis*, *Trichomonas hominis*).

E. histolytica affects about 10% of the world's population or 480 million people⁹, however this infection can be as high as 25% in certain areas of underdeveloped tropical countries. About 36 million develop clinical amebiasis and about 40,000 die annually⁹ (Table I). *G. duodenalis* is the most common intestinal protozoal infection and it is found throughout temperate and tropical regions^{10,11}. Their prevalence varies between 2%-5% in developed countries and 20%-30% in developing countries. In USA and UK, giardiasis is the most commonly reported intestinal parasitic infection of man. It is estimated that about 200 million infections occur each year in Africa, Asia and Latin America.

Human cases of *Cryptosporidium* sp. have been reported in various parts of the world and the prevalence appears to be highest in the tropics. This infection has been reported in 13% of diarrhea children in India and 7.3% in Thailand^{12,13}. In temperate countries the prevalence in children varies from 1.1% in Spain¹⁴ to 1.4% in United Kingdom¹⁵ and 2.1% in France¹⁶. The majority of the infection in man is by *Cryptosporidium parvum*.

Intestinal Parasitic Infections in Malaysia

The common intestinal parasites in man in Malaysia are mainly the helminths, especially *E. vermicularis* and the STH (*A. lumbricoides*, *T. trichiura* and *N. americanus*) and the protozoans, *E. histolytica* and *G. duodenalis*.

Several studies have demonstrated a high prevalence of *A. lumbricoides*, *T. trichiura* and *N. americanus* infections in underprivileged

community such as Orang Asli children¹⁷⁻²¹, in estates²²⁻²⁵, squatter areas²⁶⁻²⁹ and poor Malay villages^{18,30-32} (Table II). *S. stercoralis* infection is not endemic in Malaysia. It usually occurs as sporadic cases. Infections with trematodes and tapeworms (*Hymenolepis nana* and *Hymenolepis diminuta*) are only reported occasionally. The prevalence of enterobiasis both in rural and urban communities in Malaysia is high³³⁻³⁶. The overall prevalence is between 40.4-57.8%.

The prevalence of amebiasis also varies from 1% - 14%^{17,22,21,37-44} (Table III). The prevalence of giardiasis varies between 2% - 19.4%, depending on the areas studied and methods used in the detection of the protozoa^{17,22,21,37,38,40,42,43,45-47} (Table IV). In Malaysia, 4.3% of children with diarrhea were found to be positive for *Cryptosporidium* sp.⁴⁸. High prevalence of cryptosporidiosis in asymptomatic carriers among HIV-positive intravenous drug users was also reported in Malaysia⁴⁹.

Risk Factors of Intestinal Parasitic Infections

With the exception of *E. vermicularis*, intestinal trematodes and cestodes, most of the common intestinal parasitic infections of man are fecal-borne infections and the transmission occurs either directly hand-to-mouth or indirectly through food and water. For protozoal infections person-to-person transmission can occur. The source of infection is mainly human (carrier or patient). Two principal factors in maintaining endemicity of intestinal parasitic infections are favorable qualities of the soil and frequent contamination of the environment by human faeces. Their transmission within the community is predominantly related to human habits with regards to eating, defecation, personal hygiene, cleanliness and level of education. Its prevalence in the community can be used as an indicator of the conditions of living, environmental sanitation as well as the socio-economic status of the community.

Environmental factors, such as water supply for domestic and personal hygiene, sanitation and housing condition⁵⁰⁻⁵³ and other factors such as socio-economic⁵¹⁻⁵⁴, demographic^{51,53} and health related behavior^{51,52,54} are known to influence this infection. In giardiasis, one study indicates that age and the presence of infected family members were the risk factors of infection⁴⁷.

Previous studies on enterobiasis indicate that it is more prevalent in over-crowded areas or families with many members^{8,34}. Two other social factors that are often believed to be related to the transmission of *E. vermicularis* are socio-economic status and level of education although the findings are controversial^{7,34,36,56}. It can be surmised that the transmission of *E. vermicularis* is determined by the cleanliness of the house and personal hygiene and not by the size of the family and socio-economic background.

Public Health Significance of Intestinal Parasitic Infections

The amount of harm caused by intestinal parasites to the health of communities depends on several factors such as species, prevalence and/or intensity of infection, the interaction between the parasites and concurrent infections, the nutritional and immunological status of the population and numerous socio-economic factors. Their significance is extremely difficult to assess because most of these infections are asymptomatic with very low morbidity and mortality. Therefore the public health significance is always measured by the prevalence, intensity of the infection and association of these infections with human nutrition, growth and development of children and work productivity in adults.

The impact of intestinal parasitic infections on nutrition, growth and development of children has been studied since the seventies. The findings however, have remained controversial. Evidence from community studies indicate that ascariasis is associated with growth impairment⁵⁷⁻⁶⁰, impaired

lactose digestion, decreased food consumption and lower plasma vitamin A⁶¹. The role of hookworm infection in causing iron deficiency anemia has been confirmed by several studies⁶². Chronic giardiasis can interfere with the growth of children by impaired nutrient digestion (fat and vitamin) and lactose intolerance. Recently, it has been shown that trichuriasis and ascariasis impair school performance and cognitive functions of children⁶³⁻⁶⁴.

Epidemiological studies of soil-transmitted helminthiases have shown that the prevalence and intensity of infection are highest among children 4-15 years of age and significantly associated with age^{20,27}. Infection occurs at a very young age (1-2 years) and the prevalence and intensity of infection rise with age, reaching maximum values at approximately 5-7 years of age and remaining constant thereafter^{20,27}. The frequency distribution of STH infections is overdispersed and highly aggregated^{20,27,65}, which means that a large proportion of the populations are not or mildly infected and only a small proportion of them are heavily infected. Predisposition to reinfection following treatment has been reported in these infections^{29,66-69}. In highly endemic areas reinfection can occur as early as 2 months post-treatment, and by 4 months almost half of the population treated become reinfected^{20,70}. By 6 months the intensity of infection was similar to pretreatment level^{65,71}.

The recent developments and findings in epidemiological research have led to the understanding of epidemiology and population dynamics of the intestinal parasites. Mathematical models for several infections have been developed based on these findings (age-related prevalence and intensity of infection for helminthiases and age-related incidence rate for protozoal infections). These models are used for devising control strategies and planning the timing and form of intervention (chemotherapy). For example in the intervention of STH infections, periodic 4-6 monthly, targeted chemotherapy is suggested as one of the strategies.

Clinical Significance of Intestinal Parasitic Infections

Although the prevalence of intestinal parasitic infections is high, the morbidity and mortality caused by these infections is very low. They are usually considered an unimportant problem. Besides that, the statistics for hospital admissions due to intestinal parasitic infections are also scarce, although the WHO estimates suggest that there may be some 3.5 million cases admitted annually with clinical disease associated with nematode infections. Intestinal parasitic infections associated with clinical disease are well documented. Ascariasis can result in often-fatal intestinal obstruction; hookworm infection can cause iron deficiency anemia; trichuriasis is associated with chronic dysentery and rectal prolapse. Amebiasis can result in dysentery and extraintestinal complications; giardiasis is associated with acute diarrhea, steatorrhea and lactose intolerance. *Cryptosporidium parvum* and *Blastocystis hominis* have been documented as the commonest opportunistic parasites causing severe enteritis and chronic diarrhea in immunocompromised people. *Cryptosporidium parvum* has also been increasingly recognised as a cause of diarrheal illness in both immunocompetent and immunocompromised people. Ascariasis was the cause of intestinal obstruction in 5-35% of all paediatric cases in a comparison of studies conducted throughout the tropics⁷². Rectal prolapse due to trichuriasis occurred in nearly 4% of children studied in the West Indies⁷³.

In Malaysia, studies on the prevalence and clinical manifestations of intestinal parasitic infections in hospitalised patients have been carried out since the sixties. Clinical features of severe trichuriasis have been described in a few studies carried out in Hospital Kuala Lumpur. These studies indicate that rectal prolapse occurred in 50% of children with severe trichuriasis⁷⁴⁻⁷⁶. Ascariasis was responsible for 42% and 41% of all acute abdominal emergencies and intestinal obstruction respectively in children 7 years and below admitted to hospital in Kuala Lumpur⁷⁷. Ascariasis

causing acute pancreatitis and acute cholecystitis has also been reported recently in patients admitted to UMMC⁷⁸. The prevalence of intestinal parasitic infections in hospitalised patients is between 21.3-43.3%^{40,79-82}. A recent study in HUKM reported a prevalence of 5.3%⁸³.

Control and Prevention of Intestinal Parasitic Infections

The most effective control program of intestinal parasitic infections is an integrated approach with community participation. The long-term objective is to reduce the prevalence, intensity of infection and severity of intestinal parasitic infections to levels at which they cease to be of public health significance. Theoretically, the infections can be controlled and prevented by improvement in environmental sanitation such as safe methods of faeces and waste disposal and provision of safe water supplies and health education on health promotion of personal and food hygiene. Such measures are usually slow to take effect, require considerable investment and need to be accompanied by social, economic and educational development. Due to constraints at the national and individual levels, control of infection using the above methods have become unrealistic besides taking a long time.

In recent years the availability of single-dose broad-spectrum anthelmintics has helped in reducing the worm burden in endemic communities. Studies have shown that periodic chemotherapy strategy has successfully lowered the intensity of *Ascaris* and hookworm infections^{69,86,87}.

In Malaysia efforts made towards the control of STH / intestinal parasitic infections are minimal compared to other health activities. There is no national policy for the prevention and control of

these infections. Instead their control is integrated in the National Environmental Sanitation Program. The objectives of this program are to educate and to increase public awareness on personal hygiene and environmental sanitation and to give anthelmintic treatment to children. The effectiveness of this program in controlling STH and other intestinal parasitic infections is still questionable.

New Challenges in Intestinal Parasitic Infections

1. Intestinal parasitic infections are still a public health concern in Malaysia. Studies from the sixties until the nineties indicate that the prevalence and intensity of these infections are still high. They also interfere with nutrition, growth and development of children. However severe clinical consequences associated with these infections are very low. To achieve optimum quality of life do we need to consider adopting a national policy for their prevention and control as what is done for malaria and filariasis?
2. Most of the epidemiological data of these infections were acquired from sporadic studies carried out in academic and research institutions. Currently there is no database on these infections at the national level. Should we include intestinal parasitic infections as one of the parameters in our National Health and Morbidity Surveys?
3. What are the implications on medical education and training of health professionals?
4. How about intestinal parasitic infections among migrants in Malaysia. Is it a problem? If so, of what magnitude?

Above are some of the questions which pose new challenges for parasitologist of the new millennium in Malaysia.

Table I : Estimated Prevalence of Major Intestinal Parasitic Infections Worldwide

Intestinal Parasitic Infections	Estimated Prevalence (million)
Ascariasis	1000
Hookworm infection	900
Trichuriasis	500-600
Strongyloidiasis	50-100
Enterobiasis	200
Amebiasis	480

Table II: Intestinal Helminthiases in Malaysia

Author & Year / Area	Prevalence (%) of Infection		
	Ascaris	Trichuris	Hookworm
Orang Asli	(17-63)	(29-92)	(24-92)
Bisseru & Aziz (1970)	48	81	95
Che Ghani & Oothuman (1991)	43	59	26
Norhayati et al., (1997)	63	92	29
Noor Hayati et al., (1998)	17	29	24
Estates	(14-73)	(36-93)	(16-55)
Sinniah et al., (1978)	34	36	16
Zahedi et al., (1980)	52	65	28
Kan (1982)	73	55	23
Kan (1989)	34	36	16
Squatters	(23-50)	(44-62)	(4-11)
Kan (1982)	23-39	44-56	5-11
Kan & Poon (1987)	47	61	4
Bundy et al., (1988)	50	62	5
Chan et al., (1992)	47	61	4
Traditional villages	(4-75)	(13-99)	(2-26)
Khairul Anuar et al., (1978)	67	55	12
Li (1990)	68-75	67-99	6-22
Che Ghani & Oothuman, (1991)	41-43	33-59	11-26
Norhayati et al., (1995)	4	13	2

Table III: Amebiasis in Malaysia

Author & Year	Prevalence (%) of Infection
Bisseru & Aziz (1970)	1.5
Dunn (1972)	5.1
Dissanaike et al., (1977)	8.7
Sinniah et al., (1978)	1.3
Nawalinski & Roundy (1978)	1.2
Hamimah et al., (1982)	2.3
Thomas & Sinniah (1982)	8.3
Sinniah (1984)	4.4
Che Ghani et al., (1987)	14.4
Noor Hayati et al., (1998)	11.5

Table IV: Giardiasis in Malaysia

Author & Year	Prevalence (%) of Infection
Bisseru & Aziz (1970)	5.6
Dunn (1972)	10.8
Dissanaike et al., (1977)	4.8
Sinniah et al., (1978)	11.3
Nawalinski & Roundy (1978)	6.0
Hamimah et al., (1982)	2.6
Sinniah (1984)	8.5
Che Ghani et al., (1987)	9.5
Noor Hayati et al., (1998)	19.4
Norhayati et al., (1998)	19.2

Table V: Changes in the prevalence of intestinal parasitic infections in patients admitted to hospital in 1981 and 2001

Intestinal Parasites	Year of Study / %	
	1981(n=198)	2001(n=300)
<i>E. histolytica</i>	3.4	2.7
<i>G. duodenalis</i>	2.8	0.7
<i>Cryptosporidium</i> sp	-	0.3
<i>B. hominis</i>	-	0.3
<i>A. lumbricoides</i>	16.2	0.7
<i>T. trichiura</i>	15.1	1.4
Hookworm	-	0.3

References

1. Warren KS, Mahmoud AAF. (Editors). *Trop Geog Med*. New York, MacGraw-Hill Book Company, 1984.
2. Salvioli L, Bundy DAP, Tomkin A. Intestinal parasitic infections. *Trans R Soc Trop Med Hyg* 1992; 86: 353-54.
3. Walsh JA, Warren KS. Selective primary health care: an interim strategy for disease control in developing countries. *New Engl J Med* 1979; 301: 967-74.
4. Pawlowski ZS. Intestinal helminthiases and human health: recent advances and future needs. *Int J Parasitol* 1987; 17(1): 159-67.
5. Peters W. Medical aspects: comments and discussion II. In: *The relevance of parasitology to human welfare today*, Taylor AER and Muller R (Editors). *Symposia of the British Society of Parasitology*, Vol 16 Oxford: Blackwell Scientific Publications. 1978.
6. Warren KS. Hookworm control. *Lancet* 1988; 2(8616): 897-98.
7. Chan TC. Enterobiasis among school children in Macau. *Southeast Asian J Trop Med Pub Hlth* 1985; 16: 549-53.
8. Haswell-Elkins MR, Elkins D, Manjula K, Michael E, Marsdon R. The distribution and abundance of *Enterobius vermicularis* in a South Indian fishing community. *Parasitol* 1987; 95: 339-54.
9. Walsh JA. Problem in recognition and diagnosis of amebiasis. Estimation of the global magnitude of morbidity and mortality. *Rev Infect Dis* 1986; 8: 228-38.
10. Farthing MJG. Diarrhoeal disease: current concepts and future challenges pathogenesis of giardiasis. *Trans R Soc Trop Med Hyg* 1989; 87: 17-21.
11. Adam RD. The biology of *Giardia* sp. *Microbiol Rev* 1991; 55: 706-32.
12. Subramanyan VR, Broadhead RL, Pal BB. Cryptosporidiosis in children of Eastern India. *Ann Trop Paed* 1989; 9: 122-25.
13. Somchai J, Kraivichian P, Kulkumthorn M. Cryptosporidiosis among orphanage children in Thailand: A one year prospective study. *Southeast Asian J Trop Med Pub Hlth* 1990; 21(3): 458-64.
14. Garcia-Rodriguez JA, Sanchez MAM, Blasco AC. The incidence of cryptosporidiosis in children: a one year prospective survey in general hospital in Spain. *Eur J Epidemiol* 1989; 5: 70-3.
15. Baxby D, Hatt CA. The incidence of cryptosporidiosis: a two-year prospective survey in children's hospital. *J Hygiene* 1986; 96: 107-11.
16. Arnaud-Battadier F, Nacéri M, Maurage C. Cryptosporidiosis in immunocompetent patients. *New Engl J Med* 1985; 313: 1019.
17. Bisseru B, Aziz AA. Intestinal parasites, eosinophilia, haemoglobin and gamma-globulin of Malay, Chinese and Indian school children. *Med J Malaysia* 1970; 25(1): 29-33.
18. Che Ghani BM, Oothuman P. Patterns of soil-transmitted helminth infection in relation to types of water supply, housing facilities and availability of latrines in rural areas of peninsular Malaysia. In: *Collected Papers on the Control of Soil-transmitted Helminthiasis Vol V*, (Editors) Yokogawa M, et al., APCO, Tokyo, 1991; 64-71.
19. Osman A, Zaleha MI. Nutritional status of women and children in Malaysia rural populations. *Asia Pacific J Clin Nut* 1995; 4: 319-24.
20. Norhayati M, Zainudin B, Mohammod CG, Oothuman, P, Azizi O, Fatmah MS. The prevalence of *Trichuris*, *Ascaris* and hookworm infections in Orang Asli children. *Southeast Asian J Trop Med Pub Hlth* 1997; 8(1): 161-68.
21. Noor Hayati MI, Sano MI, Mohammod CG, Norhayati M, Rohani AK, Halimah AS. Infeksi parasit usus pada masyarakat Orang Asli di Pos Piah, Sungai Siput (U), Perak. *Proc 2nd National Health Science Symposium* 1998; 72-78.
22. Sinniah B, Sinniah D, Manmohan S, Poon GK. Prevalence of parasitic infections in Malaysian oil palm estate workers. *Southeast Asian J Trop Med Pub Hlth* 1978; 9(2): 272-76.
23. Zahedi M, Oothuman P, Sabapathy NN, Bakar NA. Intestinal nematode infections and efficacy study of oxfantel-pyrantel pamoate among plantation workers. *Med J Malaysia* 1980; 35: 31-7.
24. Kan SP. Soil-transmitted helminthiases in Selangor. *Med J Malaysia* 1982; 37(2): 118-22.

CONTINUING MEDICAL EDUCATION

25. Kan SP. Soil-transmitted helminthiases among inhabitants of an oil palm plantation in West Malaysia. *J Trop Med Hyg* 1989; 92: 263-69.
26. Kan SP, Poon GK. Prevalence, distribution and intensity of soil-transmitted helminthiases among Malaysian children. *Pub Hlth* 1987; 101: 243-51.
27. Bundy DAP, Kan SP, Rose R. Age-related prevalence, intensity and frequency distribution of gastrointestinal helminths infection in urban slum children from Kuala Lumpur, Malaysia. *Trans R Soc Trop Med Hyg* 1988; 82: 289-94.
28. Hanjeet K, Lai PF, Ow Yang CK, Mathias RG. Soil-transmitted helminthiases in squatter population around Kuala Lumpur by ethnic distribution. *Trop Biomed* 1991; 8: 33-7.
29. Chan L, Kan SP, Bundy DAP. The effect of repeated chemotherapy on the prevalence and intensity of infection of *Ascaris lumbricoides* and *Trichuris trichiura*. *Parasitol* 1992; 104: 371-77.
30. Khairul Anuar A, Ramachandran CP, Paran TP. Parasitic disease among fishermen living on Penang Island I. Helminthiases. *Med J Malaysia* 1978; 32(4): 321-27.
31. Li CF. Hookworm infection and protein energy malnutrition: Transverse evidence from two Malaysian ecological groups. *Trop Geog Med* 1990; 42: 8-12.
32. Norhayati M, Noor Hayati MI, Oothuman P, Fatmah MS, Muzain Minudin Y, Ismail G. Prevalence of soil-transmitted helminthiases and their relation to the socio-economic status among 1-8 year old children in rural areas in Malaysia. In: *Collected Papers on the Control of Soil-transmitted Helminthiases Vol VI*, (Editors) Yokogawa M, et al., APCO, Tokyo, 1994.
33. Oothuman P, Rampal L, Omar B, Jeffery J, Rubiah M, Ismail G, Fatmah MS. The prevalence of *Enterobius vermicularis* amongst primary school children in Pulau Ketam, Selangor. *Med J Malaysia* 1989; 44: 312-16.
34. Noor Hayati MI, Rajeswari B. Epidemiology and symptomatology of enterobiasis among young children attending a community clinic in Kuala Lumpur, Peninsular Malaysia. *Trop Biomed* 1991; 8: 151-56.
35. Rahman WA. Prevalence of *Enterobius vermicularis* in man in Malaysia. *Trans R Soc Trop Med Hyg* 1991; 85: 249.
36. Norhayati M, Noor Hayati MI, Oothuman P, Azizi O, Fatmah MS, Ismail G, Muzain Minudin Y. *Enterobius vermicularis* infection among children aged 1-8 years in a rural area in Malaysia. *Southeast Asian J Trop Med Pub Hlth* 1994; 25(3): 494-97.
37. Dunn FL. Intestinal parasitism in Malayan aborigines (Orang Asli) in Peninsular Malaysia. *Bull Wld Hlth Org* 1972; 46: 99-113.
38. Dissanaikie AS, Thomas V, Kan SP, Ong HT. Studies on parasitic infection in Orang Asli (Aborigines) in peninsular Malaysia. *Med J Malaysia* 1997; 32(1): 48-55.
39. Nawalinski T, Roundy L. Intestinal parasitism in a kampung on Pulau Pangkor, West Malaysia. *Southeast Asian J Trop Med Pub Hlth* 1978; 3: 440-51.
40. Hamimah I, Zahedi M, Ainiyah AJ. The prevalence of intestinal parasites among children at the General Hospital, Kuala Lumpur, Malaysia. *Med J Malaysia* 1982; 37(4): 373-77.
41. Thomas V, Sinniah B. Seroepidemiology of amebiasis in Peninsular Malaysia. *J Trop Med Parasitol* 1982; 76(2): 147-51.
42. Sinniah B. Intestinal protozoan and helminth infections and control of soil-transmitted helminthiases in Malaysian school children. *Pub Hlth* 1984; 98: 152-56.
43. Che Ghani BM, Mohamed AM, Oothuman P. Infection with *Entamoeba histolytica* and *Giardia lamblia* in an urban slum, rural village and medical students in Peninsular Malaysia. *Trop Biomed* 1987; 4: 150-54.
44. Lai KPF. Intestinal protozoan infections in Malaysia. *Southeast Asian J Trop Med Pub Hlth* 1992; 23: 578-86.
45. Rajeswari B, Sinniah B, Hasnah H. Socioeconomic factors associated with intestinal parasites among children living in Gombak, Malaysia. *Asia Pacific J Pub Hlth* 1994; 7: 21-5.
46. Shekar KC, Prathapa S, Gurpreet K. Prevalence of giardiasis among Malaysian primary school children. *Med J Malaysia* 1996; 51: 475-89.
47. Norhayati M, Pengabeian M, Oothuman P, Fatmah MS. Prevalence and some risk factors of *Giardia duodenalis* infection in rural communities in Malaysia. *Southeast Asian J Trop Med Pub Hlth* 1998; 29(4): 735-38.

48. Mat Ludin CM, Afifi SAB, Hasenan N, Maimunah A, Khairul A. Cryptosporidiosis among children with acute gastroenteritis in paediatric ward in the General Hospital, Penang. *Southeast Asian J Trop Med Pub Hlth* 1991; 22(2): 200-2.
49. Mohamed Kamel AG, Maning N, Arulmaithan S. Cryptosporidiosis among HIV positive intravenous drug users in Malaysia. *Southeast Asian J Trop Med Pub Hlth* 1994; 25(4): 650-53.
50. Henry FJ. Environmental sanitation, infection and nutritional status of infants in rural St. Lucia, West Indies. *Trans R Soc Trop Med Hyg* 1981; 75(4): 507-13.
51. Ismail MM, Rajapakse AL, Suraweera MGW, Weerasuriya K, Amarasinghe DKC. Some socio-economic and health-related factors and soil-transmitted nematode infections: 1. Relationship to prevalence and intensity of infection. In: *Collected Papers on the Control of Soil-Transmitted Helminthiases Vol IV*, (Editors) Yokogawa M, et al., APCO, Tokyo, 1987; 23-8.
52. Ismail MM, Rajapakse AL, Suraweera MGW, Amarasinghe DKC. Some socio-economic and health-related factors and soil-transmitted nematode infections: 2. Relationship to reinfection. In: *Collected Papers on the Control of Soil-Transmitted Helminthiases Vol V*, (Editors) Yokogawa M, et al., APCO, Tokyo, 1989; 2-35.
53. Norhayati M, Oothuman P, Fatmah MS. Some risk factors of *Ascaris* and *Trichuris* infection in Malaysian Aborigine (Orang Asli). *Med J Malaysia* 1998; 53(4): 401-07.
54. Elkins DB, Haswell-Elkins MR, Anderson RM. The epidemiology and control of intestinal helminths in the Pulikat Lake region of Southern India 1. Study design and pre- and post-treatment observations on *Ascaris lumbricoides* infection. *Trans R Soc Trop Med Hyg* 1986; 80: 774-92.
55. Bundy DAP, Cooper ES, Thompson PE, Anderson RM., Didier JM. Age related changes in the prevalence and intensity of *Trichuris trichiura* in St Lucia community. *Trans R Soc Trop Med Hyg* 1987; 81: 85-9.
56. Warren KS, Mahmoud AAF. Algorithms in the diagnosis and management of exotic disease. V. Enterobiasis. *J Infect Dis* 1975; 13: 229-32.
57. Stephenson LS, Crompton DWT, Latham MC, Schulpen TWJ, Neshiem MC, Jansen AAJ. Treatment with a single dose of albendazole improves growth of Kenyan schoolchildren with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* infection. *Am J Trop Med Hyg* 1989; 41(1): 78-87.
58. Gupta MC. Effect of ascariasis upon nutritional status of children. *J Trop Ped* 1990; 36: 189-91.
59. Thein-Hlaing, Thane-Toe, Than-Saw, Myat-Lay-Kyin. A control chemotherapeutic intervention trial on relationship between *Ascaris lumbricoides* infection and malnutrition in children. *Trans R Soc Trop Med Hyg* 1991; 85: 523-28.
60. Robertson LJ, Crompton DWT, Sanjur D, Neshiem MC. *Trichuris trichiura* and the growth of primary schoolchildren in Panama. *Trans R Soc Trop Med Hyg* 1992; 86: 656-57.
61. Taren DL, Neshiem MC, Crompton DWT, Holland CV, Barbeau I, Rivera G, Sanjur D, Tiffany J, Tucker K. Contribution of ascariasis to poor nutritional status in children from Chiriqui Province, Rep of Panama. *Parasitol* 1987; 95: 603-13.
62. Schad GA, Warren KS (Editors). *Hookworm disease*. London and Ohiladelphia: Taylor & Francis Ltd 1990.
63. Noakes C, Bundy DAP. Does helminth infection affect mental processing and educational achievement? *Parasitol Today* 1994; 10: 14-18.
64. Che Ghani M. Mental function, attention span and academic performance of school children infected with *Ascaris lumbricoides* and *Trichuris trichiura* in rural Peninsular Malaysia: the impact of anthelmintic treatment. DSc Thesis, Tulane University 1995.

Intestinal Parasitic Infections In Man: A Review

Questions

1. The following intestinal parasitic infections are endemic in Malaysia:
 - A. Trichuriasis
 - B. Giardiasis
 - C. Strongyloidiasis
 - D. Amebiasis
 - E. Fasciolopsiasis

2. Presenting symptoms of intestinal parasitic infections include
 - A. diarrhea / dysentery
 - B. respiratory symptoms
 - C. dermatitis
 - D. failure to thrive
 - E. joint pain

3. Infection with the following intestinal parasites should be ruled out in AIDS patients who present with prolonged diarrhea:
 - A. *Cryptosporidium parvum*
 - B. *Strongyloides stercoralis*
 - C. Microsporidia
 - D. *Ascaris lumbricoides*
 - E. *Isospora belli*

4. The following statements are true of intestinal protozoal infections:
 - A. They are feco-oral infections.
 - B. A few species of protozoa can cause malabsorption syndrome.
 - C. They do not cause extra-intestinal lesions.
 - D. Intestinal biopsy can be used in the diagnosis.
 - E. Most antiprotozoals are effective in their treatment.

5. The following statements are true of soil-transmitted helminthiasis:
 - A. Enterobiasis is an example.
 - B. They are zoonotic infections.
 - C. Most infections are asymptomatic.
 - D. They are easily diagnosed by fecal smear examination.
 - E. Only severe infections need treatment.